



Mark Scheme (Results)

November 2020

Pearson Edexcel International GCSE
Mathematics A (4MA1)
Paper 1HR

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

November 2020

Publications Code 4MA1_1HR_2011_MS

All the material in this publication is copyright

© Pearson Education Ltd 2020

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.

Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

- **Types of mark**

- M marks: method marks
- A marks: accuracy marks
- B marks: unconditional accuracy marks (independent of M marks)

- **Abbreviations**

- cao – correct answer only
- ft – follow through
- isw – ignore subsequent working
- SC - special case
- oe – or equivalent (and appropriate)
- dep – dependent
- indep – independent
- awrt – answer which rounds to
- eeoo – each error or omission

- **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks.

If a candidate misreads a number from the question. Eg. Uses 252 instead of 255; method marks may be awarded provided the question has not been simplified. Examiners should send any instance of a suspected misread to review. If there is a choice of methods shown, mark the method that leads to the answer on the answer line; where no answer is given on the answer line, award the lowest mark from the methods shown.

If there is no answer on the answer line then check the working for an obvious answer.

- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded to another.

International GCSE Maths A (4MA1) November 2020 – Paper 1HR Mark scheme

Apart from Questions 1, 3c, 12, 13b, 19, 23 where the mark scheme states otherwise, the correct answer, unless clearly obtained by an incorrect method, should be taken to imply a correct method.

Q	Working	Answer	Mark	Notes
1	e.g. $\frac{15}{4}$		3	M1 for $3\frac{3}{4}$ expressed as an improper fraction
	e.g. $\frac{15^5}{4} \times \frac{7}{9^3}$ OR $\frac{105}{36}$ oe			M1 correct cancelling or multiplication of numerators and denominators without cancelling
	e.g. $\frac{15^5}{4} \times \frac{7}{9^3} = \frac{35}{12} = 2\frac{11}{12}$ or $\frac{15}{4} \times \frac{7}{9} = \frac{105}{36} = \frac{35}{12} = 2\frac{11}{12}$ or $\frac{15}{4} \times \frac{7}{9} = \frac{105}{36} = 2\frac{33}{36} = 2\frac{11}{12}$	shown		A1 dep on M2, for conclusion to $2\frac{11}{12}$ from correct working – either sight of the result of the multiplication e.g. $\frac{105}{36}$ oe must be seen or correct cancelling prior to the multiplication to $\frac{35}{12}$ NB: use of decimals scores no marks
				Total 3 marks

2			2	M1 Arcs on BC , AB and arcs from these points meeting or for bisector without arcs
		Correct bisector		A1 must see correct arcs
				Total 2 marks

3	(a)		h^9	1	B1
	(b)	$(-5)^2 - 4 \times -5$ oe e.g. $25 + 20$		2	M1 for a correct substitution
			45		A1
	(c)	$5x - 3 = 4(2x + 3)$ oe or $\frac{5x}{4} - \frac{3}{4} = 2x + 3$ oe		3	M1 for correctly removing the denominator, condone missing brackets
		e.g. $5x - 8x = 12 + 3$ or $-3x = 12 + 3$ or $8x - 5x = -12 - 3$ or $3x = -12 - 3$ or $-\frac{3}{4} - 3 = 2x - \frac{5x}{4}$ or $-\frac{15}{4} = \frac{3x}{4}$			M1 for a correct rearrangement with terms in x on one side and numbers on the other, allow correct rearrangement of their equation in the form $ax + b = cx + d$
			-5		A1 dep on at least M1 SCB2 for an answer of $x = -2$ coming from $5x - 3 = 8x + 3$ or $x = 5$ coming from $5x - 3 = 2x + 12$
Total 6 marks					

4	(a)		$30 < t \leq 40$	1	B1
	(b)	e.g. $5 \times 4 + 15 \times 10 + 25 \times 15 + 35 \times 25 + 45 \times 6$ (= 1690) or $20 + 150 + 375 + 875 + 270$ (= 1690)		4	M2 For correct products using midpoints (allowing one error) with intention to add. If not M2 then award M1 for products using frequency and a consistent value within the range (allowing one error) with intention to add or correct products using midpoint without addition.
		"1690" \div 60			M1 dep on M1
			28.2		A1 accept 28.1 – 28.2
Total 5 marks					

5	(a)	$8265 - 7500 (= 765)$ or $\frac{8265}{7500} (= 1.102)$		3	M1 8265 – 7500 could be embedded in another calculation.
		$\frac{"765"}{7500} \times 100$ oe or $"1.102" \times 100 - 100$ oe			M1
			10.2		A1 oe
	(b)	e.g. $31.5(0) \div (1 - 0.3)$		3	M2 for a complete method e.g. $31.5(0) \div (1 - 0.3)$
					(M1) for $31.5(0) \div (100 - 30) (= 0.45)$ or e.g. $(1 - 0.3)x = 31.5(0)$
			45		A1
					Total 6 marks

6		e.g. $a = (-3 + 47) \div 2 (= 22)$ or $\frac{11+b}{2} = -19$ ($b = -38 - 11 = -49$) or method to add 25 to -3 or method to subtract 25 from 47 or method to subtract 30 from -19 or method to subtract 60 from 11		2	M1 for a correct method to find either coordinate or one coordinate correct. Look for correct method on their diagram, if used.
			$a = 22, b = -49$		A1 both correct
					Total 2 marks

9	e.g. $1.5 \times 1.5 (= 2.25 \text{ oe})$		3	M1 for calculating the area of the square, may be seen embedded within a calculation
	e.g. $34.8 \times "2.25"$			M1 for a complete method to find the force
		78.3		A1 oe
				Total 3 marks

10	e.g. $\frac{3}{"10"} \times 80 (= 24)$ or $\frac{2}{"10"} \times 80 (= 16)$ or $\frac{5}{"10"} \times 80 (= 40)$		5	M2 for a complete method to find the number of chocolate cakes or lemon cakes or fruit cakes "10" comes from $3 + 2 + 5$ (M1 for correct use of the ratio e.g. $80 \div "10" (= 8)$)
	e.g. $"16" \times \frac{3}{4} \times 1.7(0) (= 20.4(0))$ or $"40" \times \frac{7}{8} \times 2.4(0) (= 84)$			M1 for a method to find the profit for lemon cakes or fruit cakes
	e.g. $"24" \times 2 (= 48)$ and $"16" \times \frac{3}{4} \times 1.7(0) (= 20.4(0))$ and $"40" \times \frac{7}{8} \times 2.4(0) (= 84)$			M1 for a method to find the profit for all 3 cakes
		152.4(0)		A1
				Total 5 marks

11	(a)		9, 28, 45, 63, 76, 80	1	B1	
	(b)			2	B2	for a correct cf graph with points at ends of intervals and joined with a curve or line segments If not B2 then B1 for 5 or 6 of their points (ft from a table with only one arithmetic error) at ends of intervals and joined with a curve or line segments OR for 5 or 6 points plotted correctly at ends of intervals not joined OR for 5 or 6 of their points from table plotted consistently within each interval (not at upper ends of intervals) at their correct heights and joined with a curve or line segments
	(c)	e.g. reading across from 40 and reading down		2	M1	ft reading from a cf graph provided method is shown
			35 - 38		A1	ft from their cf graph
						Total 5 marks

12	e.g. $35x + 10y = 27.5$ or $21x + 6y = 16.5$ $\frac{6x - 10y = 34}{41x} = 61.5$ $\frac{21x - 35y = 119}{41y} = -102.5$ e.g. $3x - 5\left(\frac{5.5 - 7x}{2}\right) = 17$ or $7\left(\frac{17 + 5y}{3}\right) + 2y = 5.5$ oe		4	M1 for a correct method to eliminate x or y : coefficients of x or y the same and correct operator to eliminate selected variable (condone any one arithmetic error in multiplication) or writing x or y in terms of the other variable and correctly substituting.
		$x = 1.5$ or $y = -2.5$		A1 oe, dep on M1
				M1 (dep on 1 st M1) for a correct method to find other variable by substitution of found variable into one equation or for repeating the above method to find the second variable.
		$x = 1.5$ and $y = -2.5$		A1 oe, dep on M1
				Total 4 marks

13	(a)		$15x^2 - 2x - 6$	2	B2 for correct differentiation (B1 for 2 of $15x^2$, $-2x$, -6 correct)
	(b)	e.g. " $15x^2 - 2x - 6$ " = 2 oe		4	M1 ft, for equating their dy/dx to 2
		$15x^2 - 2x - 8 (= 0)$			M1 (dep on M1) ft their three-term quadratic
		e.g. $(3x + 2)(5x - 4) (= 0)$ $x = \frac{2 \pm \sqrt{(-2)^2 - (4 \times 15 \times -8)}}{2 \times 15}$			M1 for solving their quadratic equation using any correct method - if factorising, allow brackets which expanded give 2 out of 3 terms correct (if using formula or completing the square allow one sign error and some simplification – allow as far as e.g. $\frac{2 \pm \sqrt{4 + 480}}{30}$ oe)
			$-\frac{2}{3}, \frac{4}{5}$		A1 oe, dep on M2 (allow -0.66 or better), Both values – isw any attempt to find y coordinates
					Total 6 marks

14	$(4x+1)(x-3) = 4x^2 - 12x + x - 3 (= 4x^2 - 11x - 3)$ $(4x+1)(5x+6) = 20x^2 + 24x + 5x + 6 (= 20x^2 + 29x + 6)$ $(x-3)(5x+6) = 5x^2 + 6x - 15x - 18 (= 5x^2 - 9x - 18)$		3	M1 for multiplying 2 brackets with at least 3 out of 4 terms correct
	$(5x+6)(4x^2 - 11x - 3) = 20x^3 - 55x^2 - 15x + 24x^2 - 66x - 18$ $(x-3)(20x^2 + 29x + 6) = 20x^3 + 29x^2 + 6x - 60x^2 - 87x - 18$ $(4x+1)(5x^2 - 9x - 18) = 20x^3 - 36x^2 - 72x + 5x^2 - 9x - 18$			M1 (dep) for multiplying the product of the first 2 brackets (ft from the 1 st stage) by the 3 rd bracket, and getting at least 3 out of 6 or 4 out of 8 terms correct
		$20x^3 - 31x^2 - 81x - 18$		A1
	Alternative			
	$20x^3 + 24x^2 - 60x^2 + 5x^2 - 15x + 6x - 72x - 18$			B2 for at least 6 out of 8 terms correct (B1 for 4 or 5 out of 8 correct terms)
		$20x^3 - 31x^2 - 81x - 18$		A1
				Total 3 marks

15		$BDF = 70^\circ$	4	B1 may be marked on diagram
		<u>Alternate segment</u> theorem		B1 reason, the angle between a tangent and a chord is equal to the angle subtended in the <u>alternate segment</u>
		$EFB = 180 - (70 + 40) = 70$ <u>opposite angles</u> in a <u>cyclic quadrilateral</u>		B1 Angle EFB with reason, <u>opposite angles</u> in a <u>cyclic quadrilateral</u> sum to 180°
		$CBF = EFB$ <u>alternate</u> angles therefore EF is parallel to ABC		B1 conclusion, <u>alternate</u> angles are equal
				Total 4 marks

16	(a)		-4	1	B1
	(b)	$(f(2.6) =) 5 \times 2.6 - 7 (= 6)$ or $gf(x) = \frac{5(5x-7)}{5x-7+4}$ oe		2	M1 for finding $f(2.6)$ or $gf(x)$
			3		A1
	(c)	$5\left(\frac{5x}{x+4}\right) - 7 = 2$ or $\frac{5x}{x+4} = \frac{2+7}{5}$ oe		3	M1
		$25x = 9(x+4)$ oe			M1 for removing the denominator $(x+4)$ in a correct equation
			2.25		A1 oe
	ALT	$fg(x) = 2 \Rightarrow g(x) = f^{-1}(2) (=9/5)$ and attempt at f^{-1}			M1
	(c)	or $f^{-1}(2)$			
		$x = g^{-1}("9/5")$			M1
			2.25		A1 oe
	(d)	$y = \frac{5x}{x+4}$ or $x = \frac{5y}{y+4}$ $y(x+4) = 5x$ $x(y+4) = 5y$		3	M1
		e.g. $4y = x(5-y)$ or e.g. $4x = y(5-x)$			M1 for a correct rearrangement and factorising
			$\frac{4x}{5-x}$		A1 oe e.g. $\frac{-4x}{x-5}$
					Total 9 marks

17	(a)	$(FH =) \sqrt{12^2 + 12^2} (=16.97... \text{ or } \sqrt{288} \text{ or } 12\sqrt{2})$		3	M1
		$\tan CFH = \frac{10}{"16.97..."}$ oe or e.g. $(CF =) \sqrt{"16.97"{}^2 + 10^2} (=19.69... \text{ or } \sqrt{388} \text{ or } 2\sqrt{97})$ and e.g. $\frac{\sin CFH}{10} = \frac{\sin 90}{"19.69"}$			M1 for a correct trig statement involving CFH
			30.5		A1 accept 30.4 – 30.7
	(b)	$(BG =) 10 + \sqrt{15^2 - 12^2} (=19)$		3	M1
		$(BE =) \sqrt{"19"{}^2 + "16.97..."{}^2}$ oe			M1 ft their FH
			25.5		A1 accept 25.4 – 25.6
Total 6 marks					

18		$a + 5d = 39$ or $a + 18d = 7.8$ or $13d = -31.2$ oe		4	M1
		$a = 51$ or $d = -2.4$			A1
		e.g. $\frac{25}{2}(2 \times 51 + (25 - 1) \times -2.4)$ oe or $12.5(2a + 23d + d) = 12.5(39 + 7.8 - 2.4)$ oe			M1 for substituting their values for a and d into S_n , a and d must be clearly stated.
			555		A1
Total 4 marks					

19	8.35 or 7.25 or 6.15 or 5.25		3	B1
	$(8.35 \times 7.25) - (6.15 \times 5.25)$			M1 Allow $UB_{AD} \times UB_{DC} - LB_{EH} \times LB_{HG}$ where $8.3 < UB_{AD} \leq 8.35$, $7.2 < UB_{DC} \leq 7.25$ $6.15 \leq LB_{EH} < 6.2$, $5.25 \leq LB_{HG} < 5.3$
		28.25		A1 oe, dep on M1
				Total 3 marks

20	(i)	$(-3, -2)$	1	B1
	(ii)	$(-1.5, 4)$	1	B1 oe
				Total 2 marks

21	$12^2 = 2^4 \times 3^2$ or $2 \times 12^2 = 2^5 \times 3^2$ oe or $\frac{2 \times 12^2}{3^2} (= 32) = 2^5$		5	M1
	$18^{4n} = (2 \times 3^2)^{4n}$ or $2^{4n} \times 3^{2 \times 4n}$			M1
	$3n^2 - 14n - 5 (= 0)$			A1
	e.g. $(3n + 1)(n - 5) (= 0)$ $n = \frac{14 \pm \sqrt{(-14)^2 - (4 \times 3 \times -5)}}{2 \times 3}$			M1 for solving their 3 term quadratic equation using any correct method - if factorising, allow brackets which expanded give 2 out of 3 terms correct (if using formula or completing the square allow one sign error and some simplification – allow as far as e.g. $\frac{14 \pm \sqrt{196 + 60}}{6}$ oe)
		$-\frac{1}{3}, 5$		A1 Allow -0.33 or better for $-\frac{1}{3}$
				Total 5 marks

22	Ext angle of octagon = $360 \div 8 (= 45)$ or Int angle of octagon $(8 - 2) \times 180 \div 8$ oe (= 135)		6	M1 for method to find the size of one exterior or one interior angle of a regular octagon
	e.g. $10 + 2 \times 10 \times \sin 45 (= 10 + 10\sqrt{2}$ or 24.1...) or e.g. $\frac{10 \sin 112.5}{\sin 22.5} (= 24.1...)$			M1 method to find <i>HE</i> or <i>AD</i> 22.5 comes from $(180 - "135") \div 2$ 112.5 comes from "135" - "22.5"
	e.g. $10 \times ("10 + 10\sqrt{2} ") (= 100 + 100\sqrt{2}$ or 241.4...) or $10 \times "24.1..." (= 241.4...)$			M1 area <i>ADEH</i>
	e.g. $10 \times \sin 45^\circ (= 5\sqrt{2}$ or 7.07...) or e.g. $\sqrt{10^2 + 10^2 - 2 \times 10 \times 10 \times \cos "135"} (= 18.4...)$ or $\frac{10 \sin "135"}{\sin 22.5} (= 18.4...)$			M1 finds perpendicular height of triangle <i>ACD</i> (may be found before, but must realise this is also height of triangle) or finds the length of <i>AC</i> 22.5 comes from $(180 - "135") \div 2$
	e.g. $0.5 \times "24.1..." \times "7.07..." (= 85.3...)$ or $0.5 \times 10 \times "18.4..." \times \sin 112.5 (= 85.3...)$			M1 finds the area of triangle <i>ACD</i> 112.5 comes from "135" - "22.5"
		327		A1 accept 326 - 327
	Alternative (splitting octagon into triangles and subtracting trapezium and triangle)			
	Ext angle of octagon = $360 \div 8 (= 45)$ or Int angle of octagon $(8 - 2) \times 180 \div 8$ oe (= 135) or one of 8 angles at centre = $360 \div 8 (= 45)$		6	M1 for method to find the size of one exterior or one interior angle of a regular octagon or method to find one angle at centre of octagon when split into 8 equal triangles
	e.g. $0.5 \times 10 \times 5 \times \tan 67.5 (= 60.35...)$ or $0.5 \times \left(\frac{10 \sin 67.5}{\sin 45} \right)^2 \times \sin 45 (= 60.35...)$ or Octagon = $8 \times "60.35" (= 482.8...)$			M1 Area of one triangle (one-eighth of octagon) or octagon
	e.g. $10 + 2 \times 10 \times \sin 45^\circ (= 10 + 10\sqrt{2} = 24.14...)$			M1 Method to find <i>HE</i>
	$0.5 \times (10 + 10 + 10\sqrt{2}) \times 5\sqrt{2} (= 120.71...)$			M1 Method to find area of trapezium <i>HEGF</i>
	$0.5 \times 10 \times 10 \times \sin 135^\circ (= 35.35...)$			M1 Method to find area of triangle <i>ABC</i>
		327		A1 accept 326 - 327
Total 6 marks				

23	e.g. $\frac{3}{x+7} \times \frac{2}{x+6} + \frac{4}{x+7} \times \frac{3}{x+6} + \frac{x}{x+7} \times \frac{x-1}{x+6} (= \frac{3}{8})$ or e.g. $\frac{3}{N} \times \frac{2}{N-1} + \frac{4}{N} \times \frac{3}{N-1} + \frac{N-7}{N} \times \frac{N-8}{N-1} (= \frac{3}{8})$ oe		4	M2 for all correct products and intention to add (M1 for one correct product)
	$5x^2 - 47x + 18 = 0$ oe $(x = 9)$ or $5N^2 - 117N + 592 = 0$			M1 Correct quadratic equation
		16		A1 dep on M3
				Total 4 marks

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom